

Practice, Learn and Achieve Your Goal with Prepp

NDA Exam

Study Material for Maths

Simplifying **Government Exams**



CONTINUITY AND DIFFERENTIABILITY

Definition of Continuity

The continuity of a real function (f) on a subset of the real numbers is defined when the function exists at point c and is given as-

$$\lim_{x\to c} f(x) = f(c)$$

A real function (f) is said to be continuous if it is continuous at every point in the domain of f.

Consider a function f(x), and the function is said to be continuous at every point in [a, b] including the endpoints a and b.

Continuity of "f" at a means,

$$\lim_{x\to a} f(x) = f(a)$$

Continuity of "f" at b means,

$$\lim_{x \to b} f(x) = f(b)$$

Differentiability Formula

Assume that if f is a real function and c is a point in its domain. The derivative of f at c is defined by 0

The derivative of a function f at c is defined by-

$$\lim_{h\to 0} \frac{f(x+h) - f(c)}{h}$$

Theorem 1: Algebra of continuous functions:

If the two real functions, say f and g, are continuous at a real number c, then

(i) f + g is continuous at x=c.

(ii) f – g is continuous at x=c.

(iii) f. g is continuous at x=c.

(iv)f/g is continuous at x=c, (provided g(c) \neq 0).

Theorem 2: Suppose f and g are real-valued functions such that (f o g) is defined at c. If g is continuous at c and if f is continuous at g (c), then (f o g) is continuous at c.

Theorem 3: If a function f is differentiable at a point c, then it is also continuous at that point.

Theorem 4 (Chain Rule): Let f be a real-valued function which is a composite of two functions u and v; i.e., f = v o u.

Suppose t = u(x) and if both dt/dx and dv/dt exist, we have df/dx = (dv/dt). (dt/dx)

Theorem 5:

- (i) The derivative of e^x with respect to x is e^x ; i.e., $d/dx(e^x) = e^x$.
- (ii) The derivative of log x with respect to x is 1/x. i.e., $d/dx(\log x) = 1/x$.

Theorem 6 (Rolle's Theorem): Let $f : [a, b] \to R$ be continuous on [a, b] and differentiable on (a, b), such that f(a) = f(b), where a and b are some real numbers. Then there exists some c in (a, b) such that f'(c) = 0.

Solved Examples

Example 1: Let [.] denotes the greatest integer function and $f(x) = [tan^2x]$, then does the limit exist or is the function differentiable or continuous at 0?

```
Solution: Given f(x) = [tan^2x]

Now, -45^\circ < x < 45^\circ

tan(-45^\circ) < tan x < tan 45^\circ

-tan 45^\circ < tan x < tan 45^\circ

-1 < tan x < 1

So, 0 < tan^2x < 1

[tan^2x] = 0

So, f(x) is zero for all values of x form x = -45^\circ to 45^\circ.

Hence, f is continuous at x = 0 and f is also differentiable at 0 and has a value zero.
```

Example 2: A function is defined as follows:

$$f(x) = x^3, x^2 < 1$$

 $x, x^2 \ge 1$

Discuss the differentiability of the function at x=1.

```
Solution: We have R.H.D. = Rf'(1) = \lim_{h\to 0} (f(1-h)-f(1))/h = \lim_{h\to 0} (1+h-1)/h = 1 and L.H.D. = Lf'(1) = \lim_{h\to 0} (f(1-h)-f(1))/(-h)
```

=
$$\lim_{h\to 0} ((1-h)^3-1)/(-h)$$

$$= \lim_{h\to 0} (3-3h+h^2) = 3$$

?Rf'(1) \neq Lf'(1) \Rightarrow f(x) is not differentiable at x=1.

Example 3: If $y = (\sin^{-1}x)^2 + k \sin^{-1}x$, show that $(1-x^2) (d^2y)/dx^2 - x dy/dx = 2$

Solution: Here $y = (\sin^{-1}x)^2 + k \sin^{-1}x$.

Differentiating both sides with respect to x, we have

$$dy/dx = 2(\sin^{-1} x)/\sqrt{(1-x^2)} + k/\sqrt{(1-x^2)}$$

$$\Rightarrow$$
 (1-x²) (dy/dx)² = 4y + k²

Differentiating this with respect to x, we get

$$(1-x^2)$$
 2 dy/dx. $(d^2 y)/(dx^2)$ - 2x $(dy/dx)^2$ = 4 (dy/dx)

$$\Rightarrow$$
 (1-x²) (d²y)/dx²-x dy/dx = 2

Prepp

Latest Sarkari jobs, Govt Exam alerts, Results and Vacancies

- Latest News and Notification
- Exam Paper Analysis
- ► Topic-wise weightage
- Previous Year Papers with Answer Key
- Preparation Strategy & Subject-wise Books

To know more Click Here











